

Write your name here	
Surname	Other names
Pearson Edexcel Level 3 GCE	<div style="display: flex; justify-content: space-around;"> <div style="text-align: center;"> Centre Number <div style="border: 1px solid black; width: 40px; height: 20px; margin: 2px;"></div> <div style="border: 1px solid black; width: 40px; height: 20px; margin: 2px;"></div> <div style="border: 1px solid black; width: 40px; height: 20px; margin: 2px;"></div> <div style="border: 1px solid black; width: 40px; height: 20px; margin: 2px;"></div> <div style="border: 1px solid black; width: 40px; height: 20px; margin: 2px;"></div> </div> <div style="text-align: center;"> Candidate Number <div style="border: 1px solid black; width: 40px; height: 20px; margin: 2px;"></div> <div style="border: 1px solid black; width: 40px; height: 20px; margin: 2px;"></div> <div style="border: 1px solid black; width: 40px; height: 20px; margin: 2px;"></div> <div style="border: 1px solid black; width: 40px; height: 20px; margin: 2px;"></div> <div style="border: 1px solid black; width: 40px; height: 20px; margin: 2px;"></div> </div> </div>
<h1 style="margin: 0;">Further Mathematics</h1> <p style="margin: 5px 0;">Advanced Subsidiary</p> <p style="margin: 5px 0;">Paper 1: Core Pure Mathematics <i>Solutions</i></p>	
Sample Assessment Material for first teaching September 2017 Time: 1 hour 40 minutes	Paper Reference 8FM0/01
You must have: Mathematical Formulae and Statistical Tables, calculator	Total Marks <div style="border: 1px solid black; width: 80px; height: 40px; margin: 0 auto;"></div>

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for algebraic manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 9 questions in this question paper. The total mark for this paper is 80.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

S54456A

©2017 Pearson Education Ltd.

1/1/1/1/



Answer ALL questions. Write your answers in the spaces provided.

1. $f(z) = z^3 + pz^2 + qz - 15$

where p and q are real constants.

Given that the equation $f(z) = 0$ has roots

$$\alpha, \frac{5}{\alpha} \text{ and } \left(\alpha + \frac{5}{\alpha} - 1\right)$$

(a) solve completely the equation $f(z) = 0$

(5)

(b) Hence find the value of p .

(2)

a) $\alpha \times \frac{5}{\alpha} \times \left(\alpha + \frac{5}{\alpha} - 1\right) = 15$

$$\alpha + \frac{5}{\alpha} - 1 = 3$$

$$\alpha^2 + 5 = 4\alpha$$

$$\alpha^2 - 4\alpha + 5 = 0$$

By calc $\alpha = 2 \pm i$

Roots are $z = 2+i, \frac{5}{2+i}, 2+i + \frac{5}{2+i} - 1$

$$z = 2+i, 2-i, 3$$

b) $\sum \alpha = -p$

$$2+i + 2-i + 3 = -p$$

$$p = -7$$

2. The plane Π passes through the point A and is perpendicular to the vector \mathbf{n}

Given that

$$\overrightarrow{OA} = \begin{pmatrix} 5 \\ -3 \\ -4 \end{pmatrix} \quad \text{and} \quad \mathbf{n} = \begin{pmatrix} 3 \\ -1 \\ 2 \end{pmatrix}$$

where O is the origin,

- (a) find a Cartesian equation of Π .

(2)

With respect to the fixed origin O , the line l is given by the equation

$$\mathbf{r} = \begin{pmatrix} 7 \\ 3 \\ -2 \end{pmatrix} + \lambda \begin{pmatrix} -1 \\ -5 \\ 3 \end{pmatrix}$$

The line l intersects the plane Π at the point X .

- (b) Show that the acute angle between the plane Π and the line l is 21.2° correct to one decimal place.

(4)

- (c) Find the coordinates of the point X .

(4)

a)

$$3x - y + 2z = d$$

$$3(5) - (-3) + 2(-4) = d$$

$$15 + 3 - 8 = d$$

$$10 = d$$

Plane Π is

$$3x - y + 2z = 10$$

b)

Angle θ between line and normal to plane


$$\cos \theta = \frac{\begin{pmatrix} -1 \\ -5 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ -1 \\ 2 \end{pmatrix}}{\left| \begin{pmatrix} -1 \\ -5 \\ 3 \end{pmatrix} \right| \left| \begin{pmatrix} 3 \\ -1 \\ 2 \end{pmatrix} \right|}$$

$$= \frac{-3 + 5 + 6}{\sqrt{35} \sqrt{14}}$$

Question 2 continued

$$\theta = \cos^{-1} \left(\frac{8}{\sqrt{490}} \right) = 68.8136^\circ$$

$$\theta = 68.8^\circ$$



Angle between line and plane

$$= 90 - \theta$$

$$= 21.2^\circ$$

$$c) \quad \underline{r} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 7 - \lambda \\ 3 - 5\lambda \\ -2 + 3\lambda \end{pmatrix}$$

Sub in plane

$$3(7 - \lambda) - (3 - 5\lambda) + 2(-2 + 3\lambda) = 10$$

$$21 - 3\lambda - 3 + 5\lambda - 4 + 6\lambda = 10$$

$$8\lambda = -4$$

$$\lambda = -\frac{1}{2}$$

$$X = \left(7 - -\frac{1}{2}, 3 - 5\left(-\frac{1}{2}\right), -2 + 3\left(-\frac{1}{2}\right) \right)$$

$$X \left(\frac{15}{2}, \frac{11}{2}, -\frac{7}{2} \right)$$

(Total for Question 2 is 10 marks)

3. Tyler invested a total of £5000 across three different accounts; a savings account, a property bond account and a share dealing account.

Tyler invested £400 more in the property bond account than in the savings account.

After one year

- the savings account had increased in value by 1.5%
- the property bond account had increased in value by 3.5%
- the share dealing account had **decreased** in value by 2.5%
- the total value across Tyler's three accounts had increased by £79

Form and solve a matrix equation to find out how much money was invested by Tyler in each account.

(7)

Savings property shares
 x y z

$$x + y + z = 5000$$

$$-x + y = 400$$

$$1.015x + 1.035y + 0.975z = 5079$$

$$\text{Let } \underline{M} = \begin{pmatrix} 1 & 1 & 1 \\ -1 & 1 & 0 \\ 1.015 & 1.035 & 0.975 \end{pmatrix}$$

$$\underline{M} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 5000 \\ 400 \\ 5079 \end{pmatrix}$$

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \underline{M}^{-1} \begin{pmatrix} 5000 \\ 400 \\ 5079 \end{pmatrix} = \begin{pmatrix} 1800 \\ 2200 \\ 1000 \end{pmatrix}$$

Savings £1800 Property £2200 Shares £1000

4. The cubic equation

$$x^3 + 3x^2 - 8x + 6 = 0$$

has roots α , β and γ .

Without solving the equation, find the cubic equation whose roots are $(\alpha - 1)$, $(\beta - 1)$ and $(\gamma - 1)$, giving your answer in the form $w^3 + pw^2 + qw + r = 0$, where p , q and r are integers to be found.

(5)

$$\text{Let } w = x - 1$$

$$w + 1 = x$$

$$(w + 1)^3 + 3(w + 1)^2 - 8(w + 1) + 6 = 0$$

$$w^3 + 3w^2 + 3w + 1 + 3w^2 + 6w + 3 - 8w - 8 + 6 = 0$$

$$w^3 + 6w^2 + w + 2 = 0$$

5.

$$\mathbf{M} = \begin{pmatrix} 1 & -\sqrt{3} \\ \sqrt{3} & 1 \end{pmatrix}$$

(a) Show that \mathbf{M} is non-singular.

(2)

The hexagon R is transformed to the hexagon S by the transformation represented by the matrix \mathbf{M} .

Given that the area of hexagon R is 5 square units,

(b) find the area of hexagon S .

(1)

The matrix \mathbf{M} represents an enlargement, with centre $(0, 0)$ and scale factor k , where $k > 0$, followed by a rotation anti-clockwise through an angle θ about $(0, 0)$.

(c) Find the value of k .

(2)

(d) Find the value of θ .

(2)

$$\begin{aligned} \text{a)} \quad \det \underline{\mathbf{M}} &= 1 \times 1 - \sqrt{3} \times (-\sqrt{3}) \\ &= 1 + 3 \\ &= 4 \neq 0 \end{aligned}$$

$$\det \underline{\mathbf{M}} \neq 0 \quad \therefore \underline{\mathbf{M}} \text{ is non-singular}$$

$$\text{b)} \quad 5 \times 4 = 20 \text{ units}^2$$

$$\text{c)} \quad \begin{pmatrix} 1 & -\sqrt{3} \\ \sqrt{3} & 1 \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} k & 0 \\ 0 & k \end{pmatrix}$$

$$\begin{pmatrix} 1 & -\sqrt{3} \\ \sqrt{3} & 1 \end{pmatrix} = \begin{pmatrix} k \cos \theta & -k \sin \theta \\ k \sin \theta & k \cos \theta \end{pmatrix}$$

Question 5 continued

$$k \sin \theta = \sqrt{3}$$

$$k \cos \theta = 1$$

$$k^2 \sin^2 \theta + k^2 \cos^2 \theta = \sqrt{3}^2 + 1^2$$

$$k^2 = 4$$

$$\underline{k = 2}$$

d)

$$k \sin \theta = \sqrt{3}$$

$$2 \sin \theta = \sqrt{3}$$

$$\sin \theta = \frac{\sqrt{3}}{2}$$

$$\theta = 60^\circ \text{ or } \frac{\pi}{3}$$

(Total for Question 5 is 7 marks)

6. (a) Prove by induction that for all positive integers n ,

$$\sum_{r=1}^n r^2 = \frac{1}{6}n(n+1)(2n+1) \quad (6)$$

- (b) Use the standard results for $\sum_{r=1}^n r^3$ and $\sum_{r=1}^n r$ to show that for all positive integers n ,

$$\sum_{r=1}^n r(r+6)(r-6) = \frac{1}{4}n(n+1)(n-8)(n+9) \quad (4)$$

- (c) Hence find the value of n that satisfies

$$\sum_{r=1}^n r(r+6)(r-6) = 17 \sum_{r=1}^n r^2 \quad (5)$$

$$a) \quad \sum_{r=1}^1 r^2 = 1^2 = 1 \quad \frac{1}{6}(1)(1+1)(2+1) = 1 \quad \checkmark$$

Formula true for $n=1$

Assume true for $n=k$, then

$$\sum_{r=1}^k r^2 = \frac{1}{6}k(k+1)(2k+1)$$

$$\text{Consider } \sum_{r=1}^{k+1} r^2 = \frac{1}{6}k(k+1)(2k+1) + (k+1)^2$$

$$= \frac{1}{6}(k+1)[k(2k+1) + 6(k+1)]$$

$$= \frac{1}{6}(k+1)[2k^2 + 7k + 6]$$

$$= \frac{1}{6}(k+1)(k+2)(2k+3)$$

Question 6 continued

$$= \frac{1}{6}(k+1)((k+1)+1)(2(k+1)+1)$$

This is same formula with k replaced by $k+1$

So if formula is true for $n=k$, it is also true for $n=k+1$.

Since it is true for $n=1$, by mathematical induction it is true for all positive integers n .

$$b) \quad \sum_{r=1}^n r^3 = \frac{1}{4} n^2(n+1)^2 \quad \sum_{r=1}^n r = \frac{1}{2} n(n+1)$$

$$\begin{aligned} \sum_{r=1}^n r(r+6)(r-6) &= \sum_{r=1}^n r(r^2-36) \\ &= \sum_{r=1}^n (r^3 - 36r) \end{aligned}$$

$$= \frac{1}{4} n^2(n+1)^2 - 18n(n+1)$$

$$= \frac{1}{4} n(n+1) [n(n+1) - 72]$$

$$= \frac{1}{4} n(n+1) (n^2 + n - 72)$$

$$= \frac{1}{4} n(n+1)(n-8)(n+9)$$

(Total for Question 6 is 15 marks)

$$c) \quad \frac{1}{4} n(n+1)(n-8)(n+9) = \frac{17}{6} n(n+1)(2n+1)$$

$$\frac{1}{4} (n-8)(n+9) = \frac{17}{6} (2n+1)$$

$$n^2 + n - 72 = \frac{34}{3} (2n+1)$$

$$3n^2 + 3n - 216 = 68n + 34$$

$$3n^2 - 65n - 250 = 0$$

$$n = 25 \quad \text{or} \quad n = -\frac{10}{3}$$

$$\underline{n = 25}$$

7.

Diagrams not drawn to scale




Figure 1




Figure 2

Figure 1 shows the central cross-section $AOB CD$ of a circular bird bath, which is made of concrete. Measurements of the height and diameter of the bird bath, and the depth of the bowl of the bird bath have been taken in order to estimate the amount of concrete that was required to make this bird bath.

Using these measurements, the cross-sectional curve CD , shown in Figure 2, is modelled as a curve with equation

$$y = 1 + kx^2 \quad -0.2 \leq x \leq 0.2$$

where k is a constant and where O is the fixed origin.

The height of the bird bath measured 1.16 m and the diameter, AB , of the base of the bird bath measured 0.40 m, as shown in Figure 1.

- Suggest the maximum depth of the bird bath. (1)
 - Find the value of k . (2)
 - Hence find the volume of concrete that was required to make the bird bath according to this model. Give your answer, in m^3 , correct to 3 significant figures. (7)
 - State a limitation of the model. (1)
- It was later discovered that the volume of concrete used to make the bird bath was 0.127 m^3 correct to 3 significant figures.
- Using this information and the answer to part (c), evaluate the model, explaining your reasoning. (1)

Question 7 continued

$$y = 1 + k(0)^2 = 1 \text{ when } x = 0$$

a) Max depth = 1.16 - 1 = 0.16 m

b) $x = 0.2$ $y = 1.16$

$$y = 1 + kx^2$$

$$1.16 = 1 + 0.04k$$

$$\frac{0.16}{0.04} = k$$

$k = 4$

c) Volume = Cylinder - $\int_1^{1.16} \pi x^2 dy$

$$= \pi \times 0.2^2 \times 1.16 - \pi \int_1^{1.16} \left(\frac{y-1}{4} \right) dy$$

$$= 0.0464\pi - \frac{\pi}{4} \left[\frac{y^2}{2} - y \right]_1^{1.16}$$

$$= 0.0464\pi - \frac{\pi}{4} \left[\left(\frac{1.16^2}{2} - 1.16 \right) - \left(\frac{1}{2} - 1 \right) \right]$$

$$= 0.0464\pi - 0.0032\pi = 0.105572 \text{ m}^3$$

$= 0.136 \text{ m}^3$

(Total for Question 7 is 12 marks)

(d)	<p>Any one of e.g. the measurements may not be accurate the inside surface of the bowl may not be smooth there may be wastage of concrete when making the bird bath</p>	B1
		(1)
(e)	<p>Some comment consistent with their values. We do need a reason e.g. $\left[\left(\frac{0.136 - 0.127}{0.127} \right) \times 100 = 7.0866... \right]$ so not a good estimate because the volume of concrete needed to make the bird bath is approximately 7% lower than that predicted by the model</p> <p style="text-align: center;">or</p> <p>We might expect the actual amount of concrete to exceed that which the model predicts due to wastage, so the model does not look suitable since it predicts more concrete than was used</p>	B1ft
		(1)

8. (a) Shade on an Argand diagram the set of points

$$\left\{ z \in \mathbb{C} : |z - 4i| \leq 3 \right\} \cap \left\{ z \in \mathbb{C} : -\frac{\pi}{2} < \arg(z + 3 - 4i) \leq \frac{\pi}{4} \right\}$$


$$\arg(z - (-3 + 4i)) \quad (6)$$

The complex number w satisfies


$$|w - 4i| = 3$$

- (b) Find the maximum value of $\arg w$ in the interval $(-\pi, \pi]$.
Give your answer in radians correct to 2 decimal places.

(2)



b)



Max arg when
tangent from (0,0)
touches circle

$$\sin \alpha = \frac{3}{4}$$

$$\text{Max arg} = \frac{\pi}{2} + \sin^{-1}\left(\frac{3}{4}\right)$$

$$= 2.419$$

$$= 2.42 \text{ radians}$$

9. An octopus is able to catch any fish that swim within a distance of 2 m from the octopus's position.

A fish F swims from a point A to a point B .

The octopus is modelled as a fixed particle at the origin O .

Fish F is modelled as a particle moving in a straight line from A to B .

Relative to O , the coordinates of A are $(-3, 1, -7)$ and the coordinates of B are $(9, 4, 11)$, where the unit of distance is metres.

- (a) Use the model to determine whether or not the octopus is able to catch fish F .


(7)

- (b) Criticise the model in relation to fish F .

(1)

- (c) Criticise the model in relation to the octopus.

(1)



At closest point C

$$\vec{AB} \cdot \vec{OC} = 0$$

$$\text{Let } C \text{ be } \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \vec{OA} + \lambda \vec{AB}.$$

$$\vec{AB} = \begin{pmatrix} 12 \\ 3 \\ 18 \end{pmatrix} \quad \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -3 + 12\lambda \\ 1 + 3\lambda \\ -7 + 18\lambda \end{pmatrix}$$

$$\text{At } C \quad \begin{pmatrix} -3 + 12\lambda \\ 1 + 3\lambda \\ -7 + 18\lambda \end{pmatrix} \cdot \begin{pmatrix} 12 \\ 3 \\ 18 \end{pmatrix} = 0$$

$$-36 + 144\lambda + 3 + 9\lambda - 126 + 324\lambda = 0$$

$$477\lambda - 159 = 0$$

Question 9 continued

$$\lambda = \frac{159}{477} = \frac{1}{3}$$

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -3 + 12\left(\frac{1}{3}\right) \\ 1 + 3\left(\frac{1}{3}\right) \\ -7 + 18\left(\frac{1}{3}\right) \end{pmatrix}$$

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix}$$

$$|\vec{OC}| = \sqrt{1^2 + 2^2 + (-1)^2} = \sqrt{6} > 2$$

\therefore Octopus cannot reach fish

9(b)

e.g.

Fish F may not swim in an exact straight line from A to B Fish F may hit an obstacle whilst swimming from A to B Fish F may deviate his path to avoid being caught by the octopus

B1

(1)

(c)

e.g.

Octopus is effectively modelled as a particle – so we may need to look at where the octopus's mass is distributed

Octopus may during the fish F 's motion move away from its fixed location at O

B1

(1)

(Total for Question 9 is 9 marks)

TOTAL FOR PAPER IS 80 MARKS