Leave blank

2. A ball is thrown vertically upwards with speed u m s^{-1} from a point P at height h metres above the ground. The ball hits the ground 0.75 s later. The speed of the ball immediately before it hits the ground is 6.45 m s^{-1}. The ball is modelled as a particle.

(a) Show that $u = 0.9$

(3)

(b) Find the height above P to which the ball rises before it starts to fall towards the ground again.

(2)

(c) Find the value of h.

(3)

Leave blank

1. At time $t = 0$ a ball is projected vertically upwards from a point O and rises to a maximum height of 40 m above O. The ball is modelled as a particle moving freely under gravity.

(a) Show that the speed of projection is 28 m s^{-1}.

(3)

(b) Find the times, in seconds, when the ball is 33.6 m above O.

(5)

Leave blank

5. A stone is projected vertically upwards from a point A with speed $u\,\mathrm{m\,s^{-1}}$. After projection the stone moves freely under gravity until it returns to A. The time between the instant that the stone is projected and the instant that it returns to A is $3\frac{4}{7}$ seconds.

Modelling the stone as a particle,

(a) show that $u = 17\frac{1}{2}$,

(3)

(b) find the greatest height above A reached by the stone,

(2)

(c) find the length of time for which the stone is at least $6\frac{3}{5}$ m above A.

(6)

Leave blank

5. A particle P is projected vertically upwards from a point A with speed u m s^{-1}. The point A is 17.5 m above horizontal ground. The particle P moves freely under gravity until it reaches the ground with speed 28 m s^{-1}.

(a) Show that $u = 21$

(3)

At time t seconds after projection, P is 19 m above A.

(b) Find the possible values of t.

(5)

The ground is soft and, after P reaches the ground, P sinks vertically downwards into the ground before coming to rest. The mass of P is 4 kg and the ground is assumed to exert a constant resistive force of magnitude 5000 N on P.

(c) Find the vertical distance that P sinks into the ground before coming to rest.

(4)

Leave blank

4. The points P and Q are at the same height h metres above horizontal ground. A small stone is dropped from rest from P. Half a second later a second small stone is thrown vertically downwards from Q with speed $7.35\,\text{m}\,\text{s}^{-1}$. Given that the stones hit the ground at the same time, find the value of h.

(7)