Write your name here Surname	Other na	ames
Pearson Edexcel Level 3 GCE	Centre Number	Candidate Number
Further M Advanced Subsidiary Further Mathematics of Paper 2E: Further State	options	
Sample Assessment Material for first t Time: 1 hour 40 minutes	eaching September 2017	Paper Reference 8FM0/2E

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for algebraic manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

### Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- There are **two** sections in this question paper. Answer **all** the questions in Section A and **all** the questions in Section B.
- Answer the questions in the spaces provided
   there may be more space than you need.
- You should show sufficient working to make your methods clear.
   Answers without working may not gain full credit.
- Answers should be given to three significant figures unless otherwise stated.

#### Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 80.
- The marks for **each** question are shown in brackets
  - use this as a guide as to how much time to spend on each question.

#### Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶

S58531A
©2017 Pearson Education Ltd.
1/1/1/





#### **SECTION A**

## Answer ALL questions. Write your answers in the spaces provided.

1. A university foreign language department carried out a survey of prospective students to find out which of three languages they were most interested in studying.

A random sample of 150 prospective students gave the following results.

		Language				
		French	Spanish	Mandarin		
Gender	Male	23	22	20		
	Female	38	32	15		

A test is carried out at the 1% level of significance to determine whether or not there is an association between gender and choice of language.

(a) State the null hypothesis for this test.

(1)

(b) Show that the expected frequency for females choosing Spanish is 30.6

**(1)** 

(c) Calculate the test statistic for this test, stating the expected frequencies you have used.

(3)

(d) State whether or not the null hypothesis is rejected. Justify your answer.

**(2)** 

(e) Explain whether or not the null hypothesis would be rejected if the test was carried out at the 10% level of significance.

(1)

	<b>)</b>
Question 1 continued	
	18
	18
	18
	1 🎗
	18
	1 8
	18
	1 8
	1 8
	1 🎗
	1 8
	18
	18
	1 8

**2.** The discrete random variable X has probability distribution given by

x	-1	0	1	2	3
P(X=x)	С	а	а	b	c

The random variable Y = 2 - 5X

Given that E(Y) = -4 and  $P(Y \ge -3) = 0.45$ 

(a) find the probability distribution of X.

(7)

Given also that  $E(Y^2) = 75$ 

(b) find the exact value of Var(X)

**(2)** 

(c) Find P(Y > X)

(2)

Issue 1 – August 2017 © Pearson Education Limited 2017

	1 🖇
	$\mathbb{I} \otimes$
Question 2 continued	$\parallel \otimes$
	$\mathbb{I} \otimes$
	$\mathbb{I} \otimes$
	$\mathbb{R}^{\otimes}$
	$\mathbb{R}^{\otimes}$
	$\parallel \otimes$
	$\mathbb{I} \otimes$
	$\parallel \otimes$
	$\parallel \otimes$
	$\parallel \otimes$
	$\mathbb{I} \otimes$
	$\mathbb{R}^{\times}$
	$\parallel \otimes$
	$\mathbb{I} \otimes$
	$\mathbb{I} \otimes$
	$\parallel \otimes$
	$\parallel$ $\otimes$
	$\parallel$ $\otimes$
	$\mathbb{I} \otimes$
	$\parallel \otimes$
	$\parallel \otimes$
	$\parallel \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \$
	$\parallel \otimes$
	$\parallel \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \$
	$\mathbb{I} \otimes$
	$\parallel$ $\otimes$
	$\parallel \otimes$
	$\mathbb{I} \otimes$
	$\mathbb{I} \otimes$
	$\parallel \otimes$
	$\parallel \otimes$
	$\parallel \otimes$
	$\mathbb{I} \otimes$

3.	Two car hire companies hire cars independently of each other.	
	Car Hire A hires cars at a rate of 2.6 cars per hour.	
	Car Hire <i>B</i> hires cars at a rate of 1.2 cars per hour.	
	(a) In a 1 hour period, find the probability that each company hires exactly 2 cars.	(2)
	(b) In a 1 hour period, find the probability that the total number of cars hired by the two companies is 3	
		(2)
	(c) In a 2 hour period, find the probability that the total number of cars hired by the two companies is less than 9	
	companies is less than 7	(2)
	On average, 1 in 250 new cars produced at a factory has a defect.	
	In a random sample of 600 new cars produced at the factory,	
	(d) (i) find the mean of the number of cars with a defect,	
	(ii) find the variance of the number of cars with a defect.	(2)
	(e) (i) Use a Poisson approximation to find the probability that no more than 4 of the cars in the sample have a defect.	(-)
	(ii) Give a reason to support the use of a Doisson approximation	
	(ii) Give a reason to support the use of a Poisson approximation.	(2)
	(ii) Give a reason to support the use of a Poisson approximation.	(2)
	(ii) Give a reason to support the use of a Poisson approximation.	(2)
	(ii) Give a reason to support the use of a Poisson approximation.	(2)
	(ii) Give a reason to support the use of a Poisson approximation.	(2)
	(ii) Give a reason to support the use of a Poisson approximation.	(2)
	(ii) Give a reason to support the use of a Poisson approximation.	(2)
	(II) Give a reason to support the use of a Poisson approximation.	(2)
	(II) Give a reason to support the use of a Poisson approximation.	(2)
	(ii) Give a reason to support the use of a Poisson approximation.	(2)
	(ii) Give a reason to support the use of a roisson approximation.	(2)
	(ii) Give a reason to support the use of a Poisson approximation.	
	(ii) Give a reason to support the use of a Poisson approximation.	
	(II) Give a reason to support the use of a roisson approximation.	

	<b>\</b>
Question 3 continued	
	$\parallel$
	$\parallel$ $\otimes$
	$\parallel$ $\otimes$
	1 %
	$\parallel$
	$\mathbb{I} \otimes$
	$\mathbb{I} \otimes$
	1 8
	1 🛭
	$\parallel$ $\otimes$
	1 🖇
	1 🐰
	1 🕸
	1 8
	1 8
	1 🎗
	1 8
	1 %
	$\parallel$
	$\parallel$
	$\mathbb{I} \otimes$
	1 🎗
	1 8
	1 8
	1 8
	$\mathbb{I} \otimes$
	1 8
	$\mathbb{I} \otimes$
	1 🐰
	1 8
	1 🖇
	$\parallel \otimes$

- **4.** The discrete random variable *X* follows a Poisson distribution with mean 1.4
  - (a) Write down the value of
    - (i) P(X = 1)
    - (ii)  $P(X \leq 4)$

(2)

The manager of a bank recorded the number of mortgages approved each week over a 40 week period.

Number of mortgages approved	0	1	2	3	4	5	6
Frequency	10	16	7	4	2	0	1

(b) Show that the mean number of mortgages approved over the 40 week period is 1.4

(1)

The bank manager believes that the Poisson distribution may be a good model for the number of mortgages approved each week.

She uses a Poisson distribution with a mean of 1.4 to calculate expected frequencies as follows.

Number of mortgages approved	0	1	2	3	4	5 or more
<b>Expected frequency</b>		r	9.67	4.51	1.58	S

(c) Find the value of r and the value of s giving your answers to 2 decimal places.

**(2)** 

The bank manager will test, at the 5% level of significance, whether or not the data can be modelled by a Poisson distribution.

(d) Calculate the test statistic and state the conclusion for this test. State clearly the degrees of freedom and the hypotheses used in the test.

**(6)** 

	<b>)</b>
	1 💸
Question 4 continued	1 🕸
	1 🖇
	$\parallel \otimes$
	1 &
	1 🌣
	1 8
	1 &
	1 🌣
	1 🌣
	1 8
	$1 \otimes$
	1 🎖
	1 🕸
	$\parallel$
	$1 \otimes$
	1 🏻
	1 🕸
	1 🖔
	$\parallel$
	1 8
	1 %
	1 💸
	1 &
	1 %
	$1 \otimes$
	1 8
	1 %
	1 🕸
	1 8
	1 8
	1 8
	1 💸
	$1 \otimes$
	1 🖇

## **SECTION B**

# Answer ALL questions. Write your answers in the spaces provided

Unless otherwise indicated, whenever a numerical value of g is required, take  $g = 9.8 \,\mathrm{m\,s^{-2}}$  and give your answer to either 2 significant figures or 3 significant figures.

5.	A small ball of mass 0.1 kg is dropped from a point which is 2.4 m above a horizontal floor. The ball falls freely under gravity, strikes the floor and bounces to a height of 0.6 m above the floor. The ball is modelled as a particle.	
	(a) Show that the coefficient of restitution between the ball and the floor is 0.5	(6)
	(b) Find the height reached by the ball above the floor after it bounces on the floor for the second time.	
		(3)
	(c) By considering your answer to (b), describe the subsequent motion of the ball.	(1)

7. [In this question use  $g = 10 \text{ m s}^{-2}$ ]

A jogger of mass 60 kg runs along a straight horizontal road at a constant speed of 4 m s<sup>-1</sup>. The total resistance to the motion of the jogger is modelled as a constant force of magnitude 30 N.

(a) Find the rate at which the jogger is working.

(3)

The jogger now comes to a hill which is inclined to the horizontal at an angle  $\alpha$ , where  $\sin \alpha = \frac{1}{15}$ . Because of the hill, the jogger reduces her speed to  $3 \, \mathrm{m \, s^{-1}}$  and maintains this constant speed as she runs up the hill. The total resistance to the motion of the jogger from non-gravitational forces continues to be modelled as a constant force of magnitude  $30 \, \mathrm{N}$ .

(b) Find the rate at which she has to work in order to run up the hill at  $3 \, \text{m s}^{-1}$ .

/	
1	~ l
٧.	JI

Issue 1 - August 2017 © Pearson Education Limited 2017

- **8.** A particle *P* of mass 3*m* is moving in a straight line on a smooth horizontal table. A particle *Q* of mass *m* is moving in the opposite direction to *P* along the same straight line. The particles collide directly. Immediately before the collision the speed of *P* is *u* and the speed of *Q* is 2*u*. The velocities of *P* and *Q* immediately after the collision, measured in the direction of motion of *P* before the collision, are *v* and *w* respectively. The coefficient of restitution between *P* and *Q* is *e*.
  - (a) Find an expression for v in terms of u and e.

(6)

Given that the direction of motion of *P* is changed by the collision,

(b) find the range of possible values of *e*.

**(2)** 

(c) Show that  $w = \frac{u}{4}(1 + 9e)$ .

**(2)** 

Following the collision with P, the particle Q then collides with and rebounds from a fixed vertical wall which is perpendicular to the direction of motion of Q. The coefficient of restitution between Q and the wall is f.

Given that  $e = \frac{5}{9}$ , and that P and Q collide again in the subsequent motion,

(d) find the range of possible values of f.

(6)

	1 🖇
	$\mathbb{I} \otimes$
Question 8 continued	$\parallel \otimes$
	$\mathbb{I} \otimes$
	$\mathbb{I} \otimes$
	$\mathbb{R}^{\otimes}$
	$\mathbb{R}^{\otimes}$
	$\parallel \otimes$
	$\mathbb{I} \otimes$
	$\parallel \otimes$
	$\parallel \otimes$
	$\parallel \otimes$
	$\parallel \otimes$
	$\mathbb{R}^{\times}$
	$\parallel \otimes$
	$\mathbb{I} \otimes$
	$\parallel$ $\otimes$
	$\parallel$ $\otimes$
	$\mathbb{I} \otimes$
	$\parallel \otimes$
	$\parallel \otimes$
	$\parallel \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \$
	$\parallel \otimes$
	$\parallel \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \$
	$\mathbb{I} \otimes$
	$\parallel$ $\otimes$
	$\parallel \otimes$
	$\mathbb{I} \otimes$
	$\mathbb{I} \otimes$
	$\parallel \otimes$
	$\parallel \otimes$
	$\parallel \otimes$
	$\mathbb{I} \otimes$
	$\parallel \otimes$
	$\parallel \otimes$