


Section B

<p>7(i) $y = 2x \ln(1+x)$ $\Rightarrow \frac{dy}{dx} = \frac{2x}{1+x} + 2 \ln(1+x)$ When $x = 0$, $dy/dx = 0 + 2 \ln 1 = 0$ \Rightarrow origin is a stationary point.</p>	M1 B1 A1 E1 [4]	product rule $d/dx(\ln(1+x)) = 1/(1+x)$ soi www (i.e. from correct derivative)
<p>(ii) $\frac{d^2y}{dx^2} = \frac{(1+x).2 - 2x.1}{(1+x)^2} + \frac{2}{1+x}$ $= \frac{2}{(1+x)^2} + \frac{2}{1+x}$ When $x = 0$, $d^2y/dx^2 = 2 + 2 = 4 > 0$ $\Rightarrow (0, 0)$ is a min point</p>	M1 A1ft A1 M1 E1 [5]	Quotient or product rule on their $2x/(1+x)$ correctly applied to their $2x/(1+x)$ o.e., e.g. $\frac{4+2x}{(1+x)^2}$ cao substituting $x = 0$ into their d^2y/dx^2 www – dep previous A1
<p>(iii) Let $u = 1+x \Rightarrow du = dx$ $\Rightarrow \int \frac{x^2}{1+x} dx = \int \frac{(u-1)^2}{u} du$ $= \int \frac{(u^2 - 2u + 1)}{u} du$ $= \int (u - 2 + \frac{1}{u}) du$ *</p> $\Rightarrow \int_0^1 \frac{x^2}{1+x} dx = \int_1^2 (u - 2 + \frac{1}{u}) du$ $= \left[\frac{1}{2}u^2 - 2u + \ln u \right]_1^2$ $= 2 - 4 + \ln 2 - (\frac{1}{2} - 2 + \ln 1)$ $= \ln 2 - \frac{1}{2}$	M1 E1 B1 B1 M1 A1 [6]	$\frac{(u-1)^2}{u}$ www (but condone du omitted except in final answer) changing limits (or substituting back for x and using 0 and 1) $\left[\frac{1}{2}u^2 - 2u + \ln u \right]$ substituting limits (consistent with u or x) cao
<p>(iv) $A = \int_0^1 2x \ln(1+x) dx$ Parts: $u = \ln(1+x)$, $du/dx = 1/(1+x)$ $dv/dx = 2x \Rightarrow v = x^2$ $= \left[x^2 \ln(1+x) \right]_0^1 - \int_0^1 \frac{x^2}{1+x} dx$ $= \ln 2 - \ln 2 + \frac{1}{2} = \frac{1}{2}$ </p>	M1 A1 M1 A1 [4]	soi substituting their $\ln 2 - \frac{1}{2}$ for $\int_0^1 \frac{x^2}{1+x} dx$ cao

<p>8 (i) Stretch in x-direction s.f. $\frac{1}{2}$ translation in y-direction 1 unit up</p>	M1 A1 M1 A1 [4]	(in either order) – allow ‘contraction’ dep ‘stretch’ allow ‘move’, ‘shift’, etc – direction can be inferred from $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ or $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ dep ‘translation’. $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ alone is M1 A0
<p>(ii) $A = \int_{-\pi/4}^{\pi/4} (1 + \sin 2x) dx$ $= \left[x - \frac{1}{2} \cos 2x \right]_{-\pi/4}^{\pi/4}$ $= \pi/4 - \frac{1}{2} \cos \pi/2 + \pi/4 + \frac{1}{2} \cos (-\pi/2)$ $= \pi/2$</p>	M1 B1 M1 A1 [4]	correct integral and limits. Condone dx missing; limits may be implied from subsequent working. substituting their limits (if zero lower limit used, must show evidence of substitution) or 1.57 or better – cao (www)
<p>(iii) $y = 1 + \sin 2x$ $\Rightarrow \frac{dy}{dx} = 2\cos 2x$ When $x = 0$, $\frac{dy}{dx} = 2$ So gradient at $(0, 1)$ on $f(x)$ is 2 \Rightarrow gradient at $(1, 0)$ on $f^{-1}(x) = \frac{1}{2}$</p>	M1 A1 A1ft B1ft [4]	differentiating – allow 1 error (but not $x + 2\cos 2x$) If 1, then must show evidence of using reciprocal, e.g. $1/1$
<p>(iv) Domain is $0 \leq x \leq 2$.</p> 	B1 M1 A1 [3]	Allow 0 to 2, but not $0 < x < 2$ or y instead of x clear attempt to reflect in $y = x$ correct domain indicated (0 to 2), and reasonable shape
<p>(v) $y = 1 + \sin 2x \ x \leftrightarrow y$ $x = 1 + \sin 2y$ $\Rightarrow \sin 2y = x - 1$ $\Rightarrow 2y = \arcsin(x - 1)$ $\Rightarrow y = \frac{1}{2} \arcsin(x - 1)$</p>	M1 A1 [2]	or $\sin 2x = y - 1$ cao